

Recapitulation:

Ehrenfeucht - Fraïssé Theorem

Duplicator wins $G_k((S_V, \vec{P}), (S_U, \vec{P}))$ iff $(S_V, \vec{P}) \equiv_{k,n} (S_U, \vec{P})$.
Duplicator can respond to choices of spoiler
cannot be distinguished by FOQS-formulas of $qd \leq k$.

Application:

$(aa)^*$ is not FOQS-definable
 \Rightarrow This reasons over all FOQS-formulas.

Proof:

If $(aa)^* = L(\varphi)$ for some φ ,

then φ has $qd(\varphi) = k$ for some $k \in \mathbb{N}$.

Choose words large enough to win game:

$G_k(a^{2^k}, a^{2^k+1})$ won by duplicator.

Thus, by Ehrenfeucht - Fraïssé Theorem

a^{2^k} and a^{2^k+1} cannot be distinguished by formula of $qd \leq k$.

Thus, cannot be distinguished by φ . \square

How to prove the existence of such a winning strategy?

Lemma:

Duplicator wins $G_k(a^{2^k}, a^{2^k+1})$.

Proof:

Establish stronger result:

For all k , duplicator wins $G_k(a^i, a^j)$ with $i, j \geq 2^{k+1}$.


Proceed by induction on k :

II: Duplicator wins $G_1(a, a^n)$ with $n \geq 1$.
 $k=1$

IS: Assume duplicator has a winning strategy for $G_k(a^{i_k}, a^{j_k})$ with $i_k, j_k \geq 2^k - 1$. Consider

$$G_{k+1}(a^{i_{k+1}}, a^{j_{k+1}}) \text{ with } i_{k+1} = 2^{(k+1)} - 1 = 2 \cdot 2^k - 1 \\ j_{k+1} \geq 2^{(k+1)} - 1$$

Then



Case 1:

Spiles picks $s \leq 2^k$ in $a^{j_{k+1}}$.

Let duplicator pick $t = s \leq 2^k$ in $a^{i_{k+1}}$.

Then

$G_k(a^{s-1}, a^{s-1})$ it won by duplicator as the words coincide.

Moreover,

$G_k(a^{i_{k+1}-s}, a^{j_{k+1}-s})$ won by duplicator by the induction hypothesis.

To see this,

$$i_{k+1} - s = 2 \cdot 2^k - 1 - s \geq 2 \cdot 2^k - 1 - 2^k = 2^k - 1. \\ (\text{Def. } i_{k+1}) \quad (\text{Def. } s)$$

Case 2:

Spiles picks $s > 2^k$. Similar.

Proof (of Ehrenfeucht-Fraïssé Theorem):

" \Rightarrow " By induction on $k \in \mathbb{N}$.

IFT: Duplicator wins 0 rounds iff $\vec{s} \rightarrow \vec{t}$ is partial isomorphism. This means for every atomic formula

$$e = P_a(x) \text{ and } e = x < y$$

we have

$$S_v, \mathcal{I}[\vec{s}/\vec{x}] \models \varphi \text{ iff } S_w, \mathcal{I}[\vec{t}/\vec{x}] \models \varphi.$$

By induction on structure of formulas,

this carries over to arbitrary quantifier-free formulas.

IS: Assume a win of duplicator on $G_k((S_v, \vec{s}), (S_w, \vec{t}))$ entails $(S_v, \vec{s}) \equiv_{k, n-1} (S_w, \vec{t})$.

Consider win of duplicator on $G_{k+1}((S_v, \vec{s}), (S_w, \vec{t}))$. Show that

$$S_v, \mathcal{I}[\vec{s}/\vec{x}] \models \varphi \text{ iff } S_w, \mathcal{I}[\vec{t}/\vec{x}] \models \varphi$$

where φ is of $gd(\varphi) = k+1$.

Case $\exists x: \varphi$ (where $gd(\varphi) = k$):

If $S_v, \mathcal{I}[\vec{s}/\vec{x}] \models \exists x: \varphi$, then there is $s \in D_v$ so that

$$S_v, \mathcal{I}[\vec{s}, s/\vec{x}, x] \models \varphi.$$

Let spoiler start with s .

By assumption that duplicator wins $G_{k+1}((S_v, \vec{s}), (S_w, \vec{t}))$, can reply by position $t \in D_w$.

The resulting game

$$G_k((S_v, s, \vec{s}), (S_w, t, \vec{t}))$$

is won by duplicator.

By the induction hypothesis $(\vec{s}, s = s, \vec{s}, \vec{t}, t = t, \vec{t})$,

$$(S_v, s, \vec{s}) \equiv_{k, n-1} (S_w, t, \vec{t}).$$

As φ has $gd(\varphi) = k$, we conclude

$$S_w, \mathcal{I}[\vec{t}, t/\vec{x}, x] \models \varphi.$$

This means, there is $t \in D_w$:

$$S_w, \mathcal{I}[\vec{t}/\vec{x}] \models \exists x: \varphi.$$

Remaining cases again by induction on structure of formulas of $gd = k+1$.

\Leftarrow Let $(S_v, \vec{s}) \equiv_{k,n} (S_w, \vec{r})$.

Construct a formula $\mathcal{C}_{v,\vec{s}}^k$ of qd k that characterizes win of $G_k((S_v, \vec{s}), -)$ from duplicator's point of view:

Let $\vec{x} = (x_1, \dots, x_n)$

(Δ) iff Duplicator wins $G_k((S_v, \vec{s}), (S_w, \vec{r}))$

$$S_v, \mathbb{I}[\vec{r}/\vec{x}] \models \mathcal{C}_{v,\vec{s}}^k,$$

for arbitrary (S_w, \vec{r}) .

Assume we constructed this formula (see below how it works).

Then duplicator wins

$$G_k((S_v, \vec{s}), (S_v, \vec{s}))$$

by copying spoiler moves.

By construction of $\mathcal{C}_{v,\vec{s}}^k$ (Δ), this means

$$S_v, \mathbb{I}[\vec{s}/\vec{x}] \models \mathcal{C}_{v,\vec{s}}^k.$$

We assume that $(S_v, \vec{r}) \equiv_{k,n} (S_w, \vec{r})$.

By definition of k -equivalence and the fact that

$$\text{qd}(\mathcal{C}_{v,\vec{s}}^k) = k,$$

we get

$$S_w, \mathbb{I}[\vec{r}/\vec{x}] \models \mathcal{C}_{v,\vec{s}}^k.$$

Again with equivalence (Δ) for $\mathcal{C}_{v,\vec{s}}^k$ we get

$$G_k((S_v, \vec{s}), (S_w, \vec{r}))$$

as required.

Key tool in construction of $\mathcal{C}_{v,\vec{s}}^k$:

Lemma on winning situations for duplicator.

Construction of $\mathcal{L}_{v, \vec{s}}^k$ by induction on k :

IT1:

$$h=0 \quad \mathcal{L}_{v, \vec{s}}^0 := \bigwedge_{s_i \in \vec{s}} P_a(x_i) \quad \wedge \bigwedge_{\substack{s_i, s_j \in \vec{s} \\ s_i < s_j}} x_i < x_j$$

IS: Assume we already constructed $\mathcal{L}_{v, \vec{s}}^k$ of $qd = k$ for $G_k((S_v, \vec{s}), -)$.

Then for $G_{k+1}((S_v, \vec{s}), -)$ we set

$$\mathcal{L}_{v, \vec{s}}^{k+1} := \underbrace{\bigwedge_{s \in D_v} \exists x: \mathcal{L}_{v, s, \vec{s}}^k(x, \vec{s})}_{(a)} \quad \wedge \quad \underbrace{\forall x: \bigvee_{s \in D_u} \mathcal{L}_{v, s, \vec{s}}^k(x, \vec{s})}_{(b)}$$

By winning lemma, this yields a winning strategy for duplicator

(a) If spoiler selects from v , then there is a position x in u with which duplicator can reply (2a in lemma)

(b) Whatever spoiler chooses from u , there is a position s in v with which duplicator can respond (2b in lemma)

Quantifier depth is $k+1$.

Note:

- Selects in v by Boolean connectives
- Selects in u by quantification.

3.2 Closure properties

- Find "subclass" of regular languages that characterises FO[\exists]-definable languages
- Algebraic characterisation (as opposed to logical)
 - ↳ Focus on closure properties.